

Convergence of symmetric trap models in the hypercube

L. R. G. Fontes^{*†} P. H. S. Lima^{†‡}

Abstract

We consider symmetric trap models in the d -dimensional hypercube whose ordered mean waiting times, seen as weights of a measure in $\mathbb{N}^*$, converge to a finite measure as $d \rightarrow \infty$, and show that the models suitably represented converge to a K process as $d \rightarrow \infty$. We then apply this result to get K processes as the scaling limits of the REM-like trap model and the Random Hopping Times dynamics for the Random Energy Model in the hypercube in time scales corresponding to the *ergodic* regime for these dynamics.

Keywords and phrases trap models, weak convergence, K-process, random energy model, scaling limit, aging

2000 Mathematics Subject Classification 60K35, 60K37, 82C44

1 Introduction

Trap models have been proposed as qualitative models exhibiting localization and aging (see [1, 2] for early references). In the mathematics literature there has recently been an interest in establishing such results for a varied class of such models (see [3, 4, 5] and references therein). In particular, it has been recognized that scaling limits play an important role in such derivations (see [3, 6, 7, 8] and references therein). It may be argued that such phenomena correspond to related phenomena exhibited by limiting models.

In this paper we consider symmetric trap models in the hypercube whose mean waiting times converge as a measure to a finite measure as the dimension diverges, and show that these models converge weakly. We then apply this result to establish the scaling limits of two dynamics in the hypercube, namely the REM-like trap model and the Random Hopping Times dynamics for the Random Energy Model, in time scales corresponding to the ergodic regime for these dynamics.

^{*}Partially supported by CNPq grants 475833/2003-1, 307978/2004-4 and 484351/2006-0, and FAPESP grant 2004/07276-2.

[†]Supported by FAPESP grant 2004/13009-7

[‡]Instituto de Matemática e Estatística, Universidade de São Paulo, Rua do Matão 1010, Cidade Universitária, 05508-090 São Paulo SP, Brasil, emails: {lrenato, plima}@ime.usp.br

1.1 The model

Let $\mathcal{H}$ denote the d -dimensional hypercube, namely $\mathcal{H}$ is the graph $(\mathcal{V}, \mathcal{E})$ with

$$\begin{aligned}\mathcal{V} &= \{0, 1\}^d, \\ \mathcal{E} &= \{(v, v') \in \mathcal{V} \times \mathcal{V} : |x - x'| = 1\},\end{aligned}$$

where $|v - v'| = \sum_{i=1}^d |v(i) - v'(i)|$ is the Hamming distance in $\mathcal{V}$.

We will consider symmetric trap models in $\mathcal{H}$, namely continuous time, space inhomogeneous, simple random walks in $\mathcal{H}$, whose transition probabilities (from each site of $\mathcal{H}$ to any of its d nearest neighbors) are uniform. Let $\gamma^d = \{\gamma_v^d, v \in \mathcal{V}\}$ denote the set of mean waiting times characterizing the model.

We will map $\mathcal{V}$ onto the set $\mathcal{D} := \{1, \dots, 2^d\}$ by enumerating $\mathcal{V}$ in decreasing order of γ^d (with an arbitrary tie breaking rule), and then consider X^d , the mapped process. Let

$$\tilde{\gamma}^d = \{\tilde{\gamma}_x^d, x \in \mathcal{D}\} \quad (1.1)$$

denote the enumeration in decreasing order of γ^d , and view it as a finite measure in $\mathbb{N}^* = \{1, 2, \dots\}$, the positive natural numbers.

We next consider a class of processes which turns out to contain limits of trap models in $\mathcal{H}$ as $d \rightarrow \infty$, as we will see below. Let $\mathcal{N} := \{(N_t^{(x)})_{t \geq 0}, x \in \mathbb{N}^*\}$ be i.i.d. Poisson processes of rate 1, with $\sigma_j^{(x)}$ the j -th event time of $N^{(x)}$, $x \in \mathbb{N}^*$, $j \geq 1$, and let $\mathcal{T} = \{T_0; T_i^{(x)} i \geq 1, x \in \mathbb{N}^*\}$ be i.i.d. exponential random variables of rate 1. $\mathcal{N}$ and $\mathcal{T}$ are assumed independent. Consider now a finite measure γ supported on $\mathbb{N}^*$, and for $y \in \bar{\mathbb{N}}^* = \mathbb{N}^* \cup \{\infty\}$ let

$$\Gamma(t) = \Gamma^y(t) = \gamma_y T_0 + \sum_{x=1}^{\infty} \gamma_x \sum_{i=1}^{N_t^{(x)}} T_i^{(x)}, \quad (1.2)$$

where, by convention, $\sum_{i=1}^0 T_i^{(x)} = 0$ for every x , and $\gamma_\infty = 0$. We define the process Y on $\bar{\mathbb{N}}^*$ starting at $y \in \bar{\mathbb{N}}^*$ as follows. For $t \geq 0$

$$Y_t = \begin{cases} y, & \text{if } 0 \leq t < \gamma(y) T_0, \\ x, & \text{if } \Gamma(\sigma_j^{(x)} -) \leq t < \Gamma(\sigma_j^{(x)}) \text{ for some } 1 \leq j < \infty, \\ \infty, & \text{otherwise.} \end{cases} \quad (1.3)$$

This process, which we here call the K process with parameter γ , was introduced and studied in [7], where it was shown to arise as limits of trap models in the complete graph with n vertices as $n \rightarrow \infty$ (see Lemma 3.11 in [7]). In the next section, we derive a similar result for the hypercube. See Theorem 1. This is our main technical result. Then, in the following section, we apply that result to get the scaling limits of the REM-like trap model and the Random Hopping Times dynamics for the REM in ergodic time scales as K processes. See Section 3.

2 Convergence to the K process

Theorem 1 *Suppose that, as $d \rightarrow \infty$, $\tilde{\gamma}^d$ converges weakly to a finite measure $\tilde{\gamma}$ supported on $\mathbb{N}^*$, and that X_0^d converges weakly to a probability measure μ on $\bar{\mathbb{N}}^*$. Then, X^d converges weakly in Skorohod space as $d \rightarrow \infty$ to a K process with parameter $\tilde{\gamma}$ and initial measure μ .*

This result extends the analysis performed in [7] for the trap model in the complete graph, with a similar approach (see Lemma 3.11 in [7] and its proof). The extra difficulty here comes from the fact that the transition probabilities in the hypercube are not uniform in the state space, as is the case in the complete graph. However, all that is indeed needed is an approximate uniform entrance law in finite sets of states. This result, a key tool used several times below, is available from [9]. We state it next, in a form suitable to our purposes, but first some notation. Let $\mathcal{X}$ denote the embedded chain of X^d and for a given fixed finite subset $\mathcal{J}$ of $\mathbb{N}^*$, let $\mathcal{T}_{\mathcal{J}}$ denote the entrance time of $\mathcal{X}$ in $\mathcal{J}$, namely,

$$\mathcal{T}_{\mathcal{J}} = \inf\{n \geq 0 : \mathcal{X}_n \in \mathcal{J}\}. \quad (2.1)$$

Proposition 2 (Corollary 1.5 [9])

$$\lim_{d \rightarrow \infty} \max_{x \notin \mathcal{J}, y \in \mathcal{J}} \left| \mathbb{P}(\mathcal{X}_{\mathcal{T}_{\mathcal{J}}} = y | \mathcal{X}_0 = x) - \frac{1}{|\mathcal{J}|} \right| = 0. \quad (2.2)$$

Here $|\cdot|$ denotes cardinality.

Remark 3 *Corollary 1.5 of [9] is actually more precise and stronger than the above statement, with error of approximation estimates, and holding for $\mathcal{J}$ depending on d in a certain manner as well.*

Remark 4 *(2.2) is the only fact about the hypercube used in the proof of Theorem 1. This result would thus hold as well for other graphs with the same property. The hypercube has nevertheless been singled out in analyses of dynamics of mean field spin glasses (see above mentioned references), and that is a reason for us to do the same here.*

2.1 Proof of Theorem 1

The strategy is to approximate X^d for d large by a trap model in the complete graph with vertex set $\mathcal{M} = \{1, \dots, M\}$ and mean waiting times $\{\tilde{\gamma}_1, \dots, \tilde{\gamma}_M\}$ for $M \leq d$ large. Let Y^M denote the latter process, and let us put $Y_0^M = Y_0 \mathbf{1}\{Y_0 \in \mathcal{M}\} + W \mathbf{1}\{Y_0 \notin \mathcal{M}\}$, with W an independent uniform random variable in $\mathcal{M}$. To accomplish the approximation, we will resort to an intermediate process, which we next describe. We start by considering $\hat{X}^d$, the trap model on $\mathcal{D}$ obtained from X^d by replacing its set of mean waiting times (see (1.1) above) by $\{\tilde{\gamma}_x, x \in \mathcal{D}\}$. The intermediate process we will consider is then $\hat{X}^d$ restricted to $\mathcal{M}$, denoted $\hat{X}^{d,M}$: this is the Markov process obtained from $\hat{X}^d$ by observing it only when it is in $\mathcal{M}$ (with time stopping for $\hat{X}^{d,M}$ when $\hat{X}^d$ is outside $\mathcal{M}$).

The approximations will be strong ones: we will couple X^d to $\hat{X}^{d,M}$ and $\hat{X}^{d,M}$ to Y^M , in the spirit of Theorem 5.2 in [7], where the approximation of Y by Y^M , needed here as the last step of the argument, was established. In particular, we also couple X_0^d to Y_0 so that the former converges almost surely to the latter as $d \rightarrow \infty$.

2.1.1 Coupling of $\hat{X}^{d,M}$ and Y^M

We first look at the embedded chains of $\hat{X}^{d,M}$ and Y^M . Let $(p_{ij}^{d,M})_{i,j \in \mathcal{M}}$ be the transition probabilities of the former chain, and let $\hat{p} = \min_{i,j \in \mathcal{M}} p_{ij}^{d,M}$. We leave it to the reader to check that there is a coupling between both chains which agrees at each step with probability at least $M\hat{p}$. We resort to such a coupling. Proposition 2 implies that

$$M\hat{p} \rightarrow 1 \quad (2.3)$$

as $d \rightarrow \infty$ for every M fixed.

Since $\hat{X}^{d,M}$ and Y^M have the same mean waiting times at each site, we can couple them in such a way that they have the same waiting times at successive visits to each site. One may also find a coupling of $\hat{X}_0^{d,M}$ and Y_0^M such that

$$\mathbb{P}(\hat{X}_0^{d,M} \neq Y_0^M) \rightarrow 0 \quad (2.4)$$

as $d \rightarrow \infty$. Resorting also to that coupling, we get the following result.

Lemma 5 *For every T and M fixed, we have*

$$\mathbb{P}(\hat{X}_t^{d,M} = Y_t^M, t \in [0, T]) \rightarrow 1 \quad (2.5)$$

as $d \rightarrow \infty$.

Proof

Let N_T denote the number of jumps of Y^M in the time interval $[0, T]$, and $0 = t_0, t_1, \dots, t_{N_T}$ the respective jump times. We conclude from the above discussion that

$$\begin{aligned} \mathbb{P}(\hat{X}_t^{d,M} = Y_t^M, t \in [0, T] | N_T = k) &= \mathbb{P}(\hat{X}_{t_i}^{d,M} = Y_{t_i}^M, i = 0, 1, \dots, k | N_T = k) \\ &\geq \mathbb{P}(\hat{X}_0^{d,M} = Y_0^M) (M\hat{p})^k, \end{aligned} \quad (2.6)$$

and the result follows from (2.3), (2.4) and dominated convergence (since $M\hat{p}$ is bounded above by 1 for all d and M). $\square$

2.1.2 Coupling of X^d and $\hat{X}^{d,M}$

We couple X^d and $\hat{X}^d$ (the latter process was defined at the beginning of the section) in the following way. Note that $\mathcal{X}$ is their common embedded chain. We then make the successive sojourn times of X^d and $\hat{X}^d$, starting from the first ones, be given by $\tilde{\gamma}_{\mathcal{X}_1}^d T_1^{\mathcal{X}_1}, \tilde{\gamma}_{\mathcal{X}_2}^d T_2^{\mathcal{X}_2}, \dots$ and $\tilde{\gamma}_{\mathcal{X}_1} \hat{T}_1^{\mathcal{X}_1}, \tilde{\gamma}_{\mathcal{X}_2} \hat{T}_2^{\mathcal{X}_2}, \dots$, respectively, where the common $T_1^x, T_2^x, \dots$, $x \in \mathbb{N}^*$, are i.i.d. rate 1 exponential random variables.

With a view towards approximating X^d and $\hat{X}^{d,M}$ strongly in Skorohod space, we introduce a *time distortion function* useful for that (see (2.8) below). For K a fixed positive integer, let $\mathcal{K}$ denote the set $\{1, \dots, K\}$, and consider the successive entrance and exit times of X^d and $\hat{X}^{d,M}$ in and out of $\mathcal{K}$ defined as follows. Let $\tau_0 = \tau_0^* = \xi_0 = \xi_0^* = 0$ and for $i \geq 1$, let

$$\tau_i = \inf\{t \geq \tau_{i-1}^* : \hat{X}_t^{d,M} \in \mathcal{K}\}, \quad \tau_i^* = \inf\{t \geq \tau_i : \hat{X}_t^{d,M} \notin \mathcal{K}\}, \quad (2.7)$$




Figure 1: Trajectories of X^d and $\hat{X}^{d,M}$.

and similarly define ξ_i and ξ_i^* , $i \geq 1$ with X^d replacing $\hat{X}^{d,M}$. See Figure 1 below.

Now for $T > 0$ fixed, let $N = \min\{i \geq 1 : \tau_i > T\}$ and define

$$\tilde{\lambda}_t = \begin{cases} \xi_j + \frac{\xi_j^* - \xi_j}{\tau_j^* - \tau_j} (t - \tau_j), & \text{if } \tau_j < t \leq \tau_j^* \text{ for some } 0 \leq j < N, \\ \xi_j^* + \frac{\xi_{j+1} - \xi_j^*}{\tau_{j+1} - \tau_j^*} (t - \tau_j^*), & \text{if } \tau_j^* < t \leq \tau_{j+1} \text{ for some } 0 \leq j < N, \\ \xi_N + (t - \tau_N), & \text{if } t \geq \tau_N. \end{cases} \quad (2.8)$$

Here and below, we interpret $0/0$ as 1. See Figure 2 below.




Figure 2: Superimposed trajectories of X^d with time distorted by $\tilde{\lambda}$ and $\hat{X}^{d,M}$ (left), and superposition of the graphs of $\tilde{\lambda}$ and the identity (right).

Remark 6 With the above definition of $\tilde{\lambda}$, we first note that $\hat{X}_t^{d,M} = X_{\tilde{\lambda}_t}^d$ whenever any of both processes is visiting $\mathcal{K}$ before time τ_N .

As part of the norm in Skorohod space, we consider the class Λ of nondecreasing Lipschitz functions mapped from $[0, \infty)$ onto $[0, \infty)$, and the following

function on Λ .

$$\phi(\lambda) = \sup_{0 \leq s < t} \left| \log \frac{\lambda_t - \lambda_s}{t - s} \right| \quad (2.9)$$

We have from (2.8) above that

$$\phi(\tilde{\lambda}) \leq \max_{1 \leq j \leq N} \left| \log \frac{\xi_j - \xi_{j-1}^*}{\tau_j - \tau_{j-1}^*} \right| \vee \max_{1 \leq j \leq N} \left| \log \frac{\xi_j^* - \xi_j}{\tau_j^* - \tau_j} \right|. \quad (2.10)$$

Below, we will consider the events A_j , $j \geq 0$, as follows.

$$A_0 = \{X_0^d \in \mathcal{K}\} \cup \{\text{there exists } 0 \leq t < \xi_1 \text{ such that } X_t^d \in \mathcal{M} \setminus \mathcal{K}\}, \quad (2.11)$$

$$A_j = \{\text{there exists } \xi_j^* \leq t < \xi_{j+1} \text{ such that } X_t^d \in \mathcal{M} \setminus \mathcal{K}\}, \quad j \geq 1. \quad (2.12)$$

It follows from Proposition 2 that for $j \geq 0$

$$\lim_{d \rightarrow \infty} \inf_{x \notin \mathcal{M}} \mathbb{P}(A_j | X_{\xi_j^*}^d = x) = 1 - K/M. \quad (2.13)$$

Notice that the probability on the left hand side of (2.13) does not depend on j ; we thus get that

$$\lim_{M \rightarrow \infty} \lim_{d \rightarrow \infty} \mathbb{P}(A_j) = 1 \text{ uniformly on } j \geq 0. \quad (2.14)$$

2.1.3 Conclusion of proof of Theorem 1

Let $D_{\mathbb{N}^*}([0, \infty))$ be the (Skorohod) space of càdlàg functions of $[0, \infty)$ to $\mathbb{N}^*$ with metric

$$\rho(f, g) := \inf_{\lambda \in \Lambda} \left[\phi(\lambda) \vee \int_0^\infty e^{-u} \rho(f, g, \lambda, u) du \right], \quad (2.15)$$

where

$$\rho(f, g, \lambda, u) := \sup_{t \geq 0} |[f(t \wedge u)]^{-1} - [g(\lambda(t) \wedge u)]^{-1}|; \quad (2.16)$$

see Section 3.5 in [10]; Λ and ϕ were defined in the paragraph of (2.9) above, and $\infty^{-1} = 0$.

It follows from Lemma 3.11 in [7] that Y^M converges weakly to Y in Skorohod space as $M \rightarrow \infty$. In order to prove Theorem 1, it thus suffices to show the following result.

Lemma 7 *With the above construction of X^d and Y^M , we have that for every $\epsilon > 0$*

$$\lim_{M \rightarrow \infty} \limsup_{d \rightarrow \infty} \mathbb{P}(\rho(X^d, Y^M) > \epsilon) = 0. \quad (2.17)$$

Proof

Given $\epsilon > 0$, let $T_\epsilon = -\log(\epsilon/2)$. Then choosing λ to be the identity, noticing that ρ in (2.16) is bounded above by 1, and using Lemma 5, we find that for every $M > 0$

$$\mathbb{P}(\rho(\hat{X}^{d,M}, Y^M) > \epsilon/2) \leq \mathbb{P}(\hat{X}_t^{d,M} \neq Y_t^M \text{ for some } t \in [0, T_\epsilon]) \rightarrow 0 \quad (2.18)$$

as $d \rightarrow \infty$. So, to establish Lemma 7, it suffices to prove Lemma 8 below. $\square$

Lemma 8 *With above construction of X^d and $\hat{X}^{d,M}$, we have that for every $\epsilon > 0$*

$$\lim_{M \rightarrow \infty} \limsup_{d \rightarrow \infty} \mathbb{P}(\rho(\hat{X}^{d,M}, X^d) > \epsilon) = 0. \quad (2.19)$$

Proof

Let $T = T_\epsilon = -\log \epsilon$, choose K such that $|x^{-1} - y^{-1}| \leq \epsilon$ for every $x, y \in \bar{\mathbb{N}}^* \setminus \mathcal{K}$, and consider λ as in (2.8) with such T and K . Then, by Remark 6 and (2.9), we see that it suffices to show that for every $\epsilon > 0$

$$\lim_{M \rightarrow \infty} \limsup_{d \rightarrow \infty} \mathbb{P} \left(\max_{1 \leq j \leq N} \left| \log \frac{\xi_j^* - \xi_j}{\tau_j^* - \tau_j} \right| > \epsilon \right) = 0, \quad (2.20)$$

and

$$\lim_{M \rightarrow \infty} \limsup_{d \rightarrow \infty} \mathbb{P} \left(\max_{1 \leq j \leq N} \left| \log \frac{\xi_j - \xi_{j-1}^*}{\tau_j - \tau_{j-1}^*} \right| > \epsilon \right) = 0. \quad (2.21)$$

Proof of (2.20)

One readily checks that

$$\max_{1 \leq j \leq N} \left| \log \frac{\xi_j^* - \xi_j}{\tau_j^* - \tau_j} \right| \leq \max_{x \in \mathcal{K}} \left| \log \frac{\tilde{\gamma}_x^d}{\tilde{\gamma}_x} \right|, \quad (2.22)$$

and (2.20) follows immediately from the assumption that $\tilde{\gamma}^d \rightarrow \tilde{\gamma}$ as $d \rightarrow \infty$.

Proof of (2.21)

Let $\tilde{\gamma}^{d,K} = \{\tilde{\gamma}_x^{d,K} := \tilde{\gamma}_x^d \wedge \tilde{\gamma}_K^d, x \in \mathcal{D}\}$, and consider the trap model $X^{d,K}$ with mean waiting times $\tilde{\gamma}^{d,K}$ coupled to X^d so that both processes have the same embedded chain $\mathcal{X}$ and the respective sojourn times are given by $\tilde{\gamma}_{\mathcal{X}_1}^{d,K} T_1^{\mathcal{X}_1}, \tilde{\gamma}_{\mathcal{X}_2}^{d,K} T_2^{\mathcal{X}_2}, \dots$ and $\tilde{\gamma}_{\mathcal{X}_1}^d T_1^{\mathcal{X}_1}, \tilde{\gamma}_{\mathcal{X}_2}^d T_2^{\mathcal{X}_2}, \dots$.

Let now $\tilde{X}^{d,K}$ denote the process $X^{d,K}$ restricted to $\mathcal{K}$ (analogously as $\hat{X}^{d,M}$), with $\tilde{\mathcal{X}}$ its embedded chain. Let N^K denote the number of jumps of $\tilde{X}^{d,K}$ up to time T . Notice that N^K is a Poisson process with rate $1/\tilde{\gamma}_K^d$ independent of $\mathcal{X}$ and of the history of X^d in the time intervals $[\xi_j^*, \xi_{j+1})$, $j \geq 0$. Thus, the probability on the left hand side of (2.21) is bounded above by

$$\mathbb{P} \left(\max_{1 \leq j \leq N^K} \left| \log \frac{U_j}{V_j} \right| > \epsilon \right) \leq \sum_{n=1}^{\infty} \sum_{j=1}^n \mathbb{P} \left(\left| \log \frac{U_j}{V_j} \right| > \epsilon \right) \mathbb{P}(N^K = n), \quad (2.23)$$

where $U_j := \xi_j - \xi_{j-1}^*$ and $V_j := \tau_j - \tau_{j-1}^*$. We now estimate the first probability on the right hand side of (2.23). We first note that from (2.14), and the fact that $\mathbb{E}(N^K)$ is finite and independent of d, M , we may insert A_j in that probability. We next write $U_j = W_j + R_j$, where R_j is the time spent by X^d in $\mathcal{D} \setminus \mathcal{M}$ during the time interval $[\xi_{j-1}^*, \xi_j)$. From the elementary inequality $|\log(x+y)| \leq |\log x| + y$, valid for all $x, y > 0$, we get that

$$\mathbb{P} \left(\left| \log \frac{U_j}{V_j} \right| > \epsilon, A_j \right) \leq \mathbb{P} \left(\left| \log \frac{W_j}{V_j} \right| > \epsilon/2, A_j \right) + \mathbb{P}(R_j > \epsilon V_j/2). \quad (2.24)$$

Arguing as in (2.22) above, we find that the first event in the first probability on the right hand side of (2.24) is empty as soon as $\max_{x \in \mathcal{M}} \left| \log \frac{\tilde{\gamma}_x^d}{\tilde{\gamma}_x} \right| \leq \epsilon/2$,

thus from $\tilde{\gamma}^d \rightarrow \tilde{\gamma}$ as $d \rightarrow \infty$ we only need to consider the second probability on the right hand side of (2.24). One readily checks that it is bounded above by

$$\max_{x \notin \mathcal{K}} \mathbb{P} \left(R_j > \epsilon V_j / 2 \mid X_{\xi_{j-1}^*}^d = x \right) \quad (2.25)$$

for all $j \geq 0$, and that the above expression does not depend on j . It is enough then to show that for any $\epsilon > 0$

$$\mathbb{P}(R_1 > \epsilon V_1 \mid X_0^d = x) =: \mathbb{P}_x(R_1 > \epsilon V_1) \rightarrow 0 \quad (2.26)$$

as $d \rightarrow \infty$ and then $M \rightarrow \infty$, uniformly in $x > K$. This is readily seen to follow from the facts that

$$\lim_{M \rightarrow \infty} \limsup_{d \rightarrow \infty} \max_{x \notin \mathcal{K}} \mathbb{P}_x(R_1 > \epsilon) = 0 \quad (2.27)$$

for any $\epsilon > 0$, and that, given $\delta > 0$, there exists $\epsilon > 0$ such that

$$\limsup_{M \rightarrow \infty} \limsup_{d \rightarrow \infty} \max_{x \notin \mathcal{K}} \mathbb{P}_x(V_1 \leq \epsilon) \leq \delta. \quad (2.28)$$

Proof of (2.27)

Let $x \notin \mathcal{K}$ be arbitrary. We will estimate

$$\mathbb{E}_x(R_1) := \mathbb{E}(R_1 \mid X_0^d = x) = \sum_{y=M+1}^d \tilde{\gamma}_y^d \mathbb{E}_x(\mathcal{L}(y)), \quad (2.29)$$

where $\mathcal{L}(y)$ is the number of visits of $\mathcal{X}$ to y from time 0 till its first entrance in $\mathcal{K}$.

Let $\mathcal{K}_y = \mathcal{K} \cup \{y\}$ and consider the discrete time Markov process $\bar{\mathcal{X}}$ such that $\bar{\mathcal{X}}_0 = x$ and otherwise $\bar{\mathcal{X}}$ is the restriction of $\mathcal{X}$ to $\mathcal{K}_y = \mathcal{K} \cup \{y\}$, and let $\bar{\mathcal{L}}(y)$ denote the number of visits of $\bar{\mathcal{X}}$ to y from time 0 till its first entrance in $\mathcal{K}$. Clearly,

$$\mathcal{L}(y) = \bar{\mathcal{L}}(y). \quad (2.30)$$

Now let $\mathcal{X}^*$ denote the Markov chain on $\mathcal{K}_y \cup \{x\}$ with the following set of transition probabilities. Let $p^1 = (p_{wz}^1, w, z \in \mathcal{K}_y \cup \{x\})$, and $p^2 = (p_{wz}^2, w, z \in \mathcal{K}_y \cup \{x\})$ denote the sets of transition probabilities of $\bar{\mathcal{X}}$ and $\mathcal{X}^*$, respectively. We make

$$p_{xy}^2 = p_{yy}^2 = p^* := \max\{p_{wz}^1; w = x, y; z \in \mathcal{K}_y\}, \quad (2.31)$$

and the remaining p_{wz}^2 can be assigned arbitrarily with the only obvious condition that p^2 is a set of transition probabilities on $\mathcal{K}_y$. Let $\mathcal{L}^*(y)$ denote the number of visits of $\mathcal{X}^*$ to y from time 0 till its first entrance in $\mathcal{K}$. One readily checks that $\mathcal{L}^*(y)$ is a Geometric random variable with parameter $1 - p^*$ and that it stochastically dominates $\bar{\mathcal{L}}(y)$. From this and (2.30), we conclude that

$$\mathbb{E}_x(\mathcal{L}(y)) \leq \frac{p^*}{1 - p^*} \quad (2.32)$$

uniformly in $x \notin \mathcal{K}$. Proposition 2 then implies that

$$\limsup_{d \rightarrow \infty} \max_{x \notin \mathcal{K}} \mathbb{E}_x(\mathcal{L}(y)) \leq \frac{1}{K}. \quad (2.33)$$

It follows readily from this, (2.29) and the assumption that $\tilde{\gamma}^d \rightarrow \tilde{\gamma}$ as $d \rightarrow \infty$ that

$$\limsup_{d \rightarrow \infty} \max_{x \notin \mathcal{K}} \mathbb{E}_x(R_1) \leq \frac{1}{K} \sum_{y=M+1}^{\infty} \tilde{\gamma}_y, \quad (2.34)$$

and (2.27) follows (using Markov's inequality), since $\tilde{\gamma}$ is a finite measure on $\mathbb{N}^*$.

Proof of (2.28)

Let us fix $x_0 \notin \mathcal{K}$. Consider the Markov process $Z = (Z_t)_{t \geq 0}$ on $\mathcal{M}$ such that $Z_0 = x_0$, for every $x \in \mathcal{M}$ the waiting time at x before jumping is exponential with mean $\tilde{\gamma}_x$, and the transition probability to $y \in \mathcal{M}$ equals $\hat{p}$, if $y \notin \mathcal{K}$, and $\frac{1-(M-K)\hat{p}}{K}$, if $y \in \mathcal{K}$, where $\hat{p}$ was defined in the paragraph of (2.3) above.

One readily checks that, given $X_0^d = x_0$, V_1 stochastically dominates S , the time Z spends in $\mathcal{M} \setminus \mathcal{K}$ before hitting $\mathcal{K}$ for the first time. Since $\tilde{\gamma}_x$ is decreasing in x , by the construction of Z , we have that, for every $L \in \{K+1, \dots, M\}$, S dominates stochastically the random variable $\tilde{\gamma}_L T 1_C$, where C is the event that Z visits $\{K+1, \dots, L\}$ before hitting $\mathcal{K}$ for the first time, T is an exponential random variable of rate 1, and $1_{\cdot}$ is the usual indicator function.

Let now $\mathcal{Z}$ denote the embedded chain for Z , and $\mathcal{T}_L = \inf\{n \geq 1 : \mathcal{Z}_n \in \{1, \dots, L\}\}$. Then

$$\begin{aligned} & \mathbb{P}(C | \mathcal{Z}_0 = x_0) \\ &= \mathbb{P}(\mathcal{Z}_{\mathcal{T}_L} \in \{K+1, \dots, L\} | \mathcal{Z}_{\mathcal{T}_L} \leq L, \mathcal{Z}_0 = x_0) \\ &= \frac{(L-K)\hat{p}}{K \left[\frac{1-(M-K)\hat{p}}{K} \right] + (L-K)\hat{p}} = \frac{(L-K)\hat{p}}{1-(M-L)\hat{p}} \\ &= \frac{(L-K)M\hat{p}}{(1-M\hat{p})M + LM\hat{p}} \rightarrow \frac{(L-K)}{L} = 1 - \frac{K}{L} \end{aligned} \quad (2.35)$$

as $d \rightarrow \infty$ uniformly in x_0 ; see (2.3).

We conclude that

$$\limsup_{M \rightarrow \infty} \limsup_{d \rightarrow \infty} \max_{x \notin \mathcal{K}} \mathbb{P}_x(V_1 \leq \epsilon) \leq \mathbb{P}(\tilde{\gamma}_L T \leq \epsilon) + \frac{K}{L} \quad (2.36)$$

for every $K < L$. Thus, given K and $\delta > 0$, we first choose L such that $\frac{K}{L} \leq \delta/2$, and then $\epsilon > 0$ such that $\mathbb{P}(\tilde{\gamma}_L T \leq \epsilon) \leq \delta/2$, and we are done. $\square$

3 The REM-like trap model and the Random hopping times dynamics for the REM

In this section we apply Theorem 1 to obtain the scaling limits of two disordered trap models in the hypercube, namely trap models in the hypercube whose mean waiting times are random variables.

3.1 The REM-like trap model

Let $\tau^d := \{\tau_v^d, v \in \mathcal{V}\}$, an i.i.d. family of random variables in the domain of attraction of an α -stable law with $0 < \alpha < 1$, be the mean waiting times of a trap model in $\mathcal{H}$. Let us then consider as before $\tilde{\tau}^d := \{\tilde{\tau}_x^d, x \in \mathcal{D}\}$, the

decreasing order statistics of τ (with an arbitrary tie breaking rule), and let Y^d be the mapped process on $\mathcal{D}$.

Now let c_d be a scaling factor such that $\hat{\tau}^d := c_d \tilde{\tau}^d$ converges to the increments in $[0, 1]$ of an α -stable subordinator. Namely,

$$c_d = \left(\inf\{t \geq 0 : \mathbb{P}(\tau_0 > t) \leq 2^{-d}\} \right)^{-1}. \quad (3.1)$$

Let us now consider Y^d speeded up by c_d , namely $\hat{Y}_t^d = Y^d(t/c_d)$, $t \geq 0$. Notice that $\hat{Y}^d$ is a trap model on $\mathcal{H}$ with mean waiting times given by $\hat{\tau}^d$.

Let $\hat{\gamma} = \{\hat{\gamma}_i, i \in \mathbb{N}^*\}$ denote the increments in $[0, 1]$ of an α -stable subordinator in decreasing order.

Corollary 9 *Suppose that $\hat{\tau}_0^d$ converges weakly to a probability measure μ on $\bar{\mathbb{N}}^*$. Then*

$$(\hat{Y}^d, \hat{\tau}^d) \Rightarrow (Y, \hat{\gamma}), \quad (3.2)$$

where Y is a K process with parameter $\hat{\gamma}$ and initial measure μ , and $\Rightarrow$ means weak convergence in the product of Skorohod norm and weak convergence norm in the space of finite measures in $\mathbb{N}^*$.

Remark 10 *In [7], a similar result was proved for the REM-like trap model in the complete graph. See Theorem 5.2 in that reference.*

Proof of Corollary 9

We can suppose that we are in a probability space where $\hat{\tau}^d \rightarrow \hat{\gamma}$ almost surely (see proof of Theorem 5.2 in [7] for an explicit argument). We can then invoke Theorem 1 to get that $\hat{Y}^d \Rightarrow Y$, and the full result follows. $\square$

3.2 Random Hopping Times dynamics for the REM

This is a dynamics whose equilibrium is the Random Energy Model. Let $H^d := \{H_v^d, v \in \mathcal{V}\}$ be an i.i.d. family of standard normal random variables, and make $\tau^d := \{\tau_v^d, v \in \mathcal{V}\}$, where $\tau_v^d = e^{\beta\sqrt{d}H_v^d}$. Defining now $\tilde{\tau}^d$, $\hat{\tau}^d$, Y^d and $\hat{Y}^d$ as above, with

$$c_d = e^{-\frac{2\log 2}{\alpha}d + \frac{1}{2\alpha}\log d}, \quad (3.1)$$

with $\alpha = \sqrt{2\log 2}/\beta$, we have that, if $\alpha < 1$, then Corollary 9 holds in this case as well, with $\hat{\gamma}$ and μ as before.

The proof starts from the known result that in this case $\hat{\tau}^d \Rightarrow \hat{\gamma}$ (see Remark to Theorem 2 in [11]). Again, as in the proof of Corollary 9 above, we can go to a probability space where the latter convergence is almost sure, and close the argument in the same way.

Remark 11 *The time scale $t \rightarrow t/c_d$ adopted in the above models is the ergodic time scale mentioned in [12]. Under shorter scalings (i.e., $t \rightarrow t/c'_d$, with $c'_d \gg c_d$) the model exhibits aging (when starting from the uniform distribution), and under longer ones ($c'_d \ll c_d$), the model reaches equilibrium. More precisely, under shorter scalings, we have that as $d \rightarrow \infty$*

$$\mathbb{P}_{\mu^d}(Y^d(t/c'_d) = Y^d((t+s)/c'_d)) \rightarrow \mathcal{R}(s/t), \quad (3.2)$$

where μ^d is the initial uniform distribution on $\mathcal{D}$, and $\mathcal{R}$ is a nontrivial function such that $\mathcal{R}(0) = 1$ and $\lim_{x \rightarrow \infty} \mathcal{R}(x) = 0$. Indeed, for the models of this section (as well as in many other instances in the references), $\mathcal{R}$ is the arcsine law:

$$\mathcal{R}(x) = \frac{\sin(\pi\alpha)}{\pi} \int_{\frac{x}{1+x}}^1 s^{-\alpha}(1-s)^{\alpha-1} ds. \quad (3.3)$$

See [13]. Under longer scalings, it can be shown that $Y^d(t/c'_d) \Rightarrow \bar{\gamma}$ as $d \rightarrow \infty$ for every $t > 0$, where $\bar{\gamma}$ is $\hat{\gamma}$ normalized to be a probability measure. It is the limiting equilibrium measure, or more precisely, the equilibrium measure of Y .

Remark 12 It can be shown that Y exhibits aging at a vanishing time scale (when starting from ∞), i.e.

$$\mathbb{P}_\infty(Y(\epsilon t) = Y(\epsilon(t+s))) \rightarrow \mathcal{R}(s/t) \quad (3.4)$$

as $\epsilon \rightarrow 0$. See Theorem 5.11 in [7]. This is in agreement with (3.2).

Acknowledgements

This work was mostly done as part of the master's project of the second author, at IME-USP and financed by FAPESP. We thank Claudio Landim for pointing out a mistake in a previous version of this paper.

References

- [1] T. M. Nieuwenhuizen; M. H. Ernst (1985) Excess noise in a hopping model for a resistor with quenched disorder, *J. Stat. Phys.* **41**, 773-801
- [2] J.-P. Bouchaud (1992) Weak ergodicity breaking and aging in disordered systems, *J. Phys. I (France)* **2**, 1705-1713
- [3] L. R. G. Fontes, M. Isopi, C. M. Newman (2002) Random walks with strongly inhomogeneous rates and singular diffusions: convergence, localization and aging in one dimension, *Ann. Probab.* **30**, 579-604
- [4] G. Ben Arous, A. Bovier, V. Gayrard (2003) Glauber dynamics of the random energy model. II. Aging below the critical temperature, *Comm. Math. Phys.* **236**, no. 1, 1-54.
- [5] G. Ben Arous, J. Černý, T. Mountford (2006) Aging in two-dimensional Bouchaud's model, *Probab. Theory Related Fields* **134**, 1-43.
- [6] G. Ben Arous, J. Černý (2007) Scaling limit for trap models on $\mathbb{Z}^d$, *Ann. Probab.* **35**, 2356-2384
- [7] L. R. G. Fontes; P. Mathieu (2008) K-processes, scaling limit and aging for the trap model in the complete graph, *Ann. Probab.* **36**, 1322-1358
- [8] G. Ben Arous, A. Bovier, J. Černý (2008) Universality of the REM for dynamics of mean-field spin glasses, *Commun. Math. Phys.* **282**, 663-695

- [9] G. Ben Arous, V. Gayrard (2008) Elementary potential theory on the hypercube, *Electron. J. Probab.* **13**, no. 59, 1726-1807
- [10] Ethier, S.N.; Kurtz, T.G. (1986) *Markov processes. Characterization and convergence*, Wiley
- [11] A. Galves; S. Martínez; P. Picco (1989) Fluctuations in Derrida's random energy and generalized random energy models, *J. Statist. Phys.* **54**, 515-529
- [12] J.-P. Bouchaud; D. S. Dean (1995) Aging on Parisi's tree, *J. Phys. I France* **5**, 265-286.
- [13] G. Ben Arous, V. Černý (2006) *Dynamics of Trap Models*, Course at the Les Houches Summer School on Mathematical Statistical Physics, Elsevier